

Content



0. Introduction

1. Differential equations

1.1 Classical equation of motion (classical mechanics, pendulum)

1.2 Partial differential equation relaxation methods (electromagnetism, diffusion)

1.3 Partial differential equation in space-time (traffic flow, tsunami)

2. Eigenvalue problem

2.1 Schrödinger equation and Hamiltonian (Harmonic oscillator, wave package)

2.2 Quantum lattice model and Hilbert space (Heisenberg model)

2.3 Exact diagonalization of spin chain (Spin wave, Haldane conjecture, topology)

2.4 Matrix product state and density matrix renormalization group (DMRG)

Content



3. Statistical and many-body physics

3.1 Classical Monte Carlo and phase transitions (Ising model and critical phenomena)

3.2 Quantum Monte Carlo methods (Path-integral and cluster update)

4. Machine learning in physics and High performance computation

4.1 AI in quantum physics

4.2 HPC and parallelism

4.3 ...

Schrödinger equation

$$i\hbar \frac{\partial}{\partial t} \psi(t, \vec{x}) = -\frac{\hbar^2}{2m} \Delta \psi(t, \vec{x}) + V(\vec{x}) \psi(t, \vec{x}) = H(\vec{x}) \psi(t, \vec{x})$$

$$\psi(t, \vec{x}) = e^{-it \frac{E}{\hbar}} \psi(\vec{x})$$


$$H(\vec{x}) \psi(\vec{x}) = E \psi(\vec{x})$$

Animating Schrödinger's Equation

<https://www.youtube.com/watch?v=Xj9PdeY64rA>


Particle in a box

https://en.wikipedia.org/wiki/Particle_in_a_box



Quantum harmonic oscillator

https://en.wikipedia.org/wiki/Quantum_harmonic_oscillator



Hydrogen atom



$$\left(-\frac{\hbar^2}{2\mu} \Delta - \frac{e^2}{4\pi\epsilon_0 r}\right) \psi(r, \theta, \phi) = E \psi(r, \theta, \phi) \quad \mu = \frac{m_e M}{(m_e + M)}$$

in spherical coordinates

$$-\frac{\hbar^2}{2\mu} \left[\frac{1}{r^2} \frac{\partial}{\partial r} \left(r^2 \frac{\partial \psi}{\partial r} \right) + \frac{1}{r^2 \sin(\theta)} \frac{\partial}{\partial \theta} \left(\sin(\theta) \frac{\partial \psi}{\partial \theta} \right) + \frac{1}{r^2 \sin^2(\theta)} \frac{\partial^2 \psi}{\partial \phi^2} \right] - \frac{e^2}{4\pi\epsilon_0 r} \psi = E \psi$$

$$\psi_{n,l,m}(r, \theta, \phi) = \sqrt{\left(\frac{2}{na_0^*}\right)^3 \frac{(n-l-1)!}{2n(n+l)!}} e^{-\rho/2} \rho^l L_{n-l-1}^{2l+1}(\rho) Y_l^m(\theta, \phi) \quad E_n = -\frac{m_e e^4}{2(4\pi\epsilon_0)^2 \hbar^2} \frac{1}{n^2}$$

$$n = 1, E_1 = 13.6 \text{ eV} = 1 \text{ Rydberg}$$



Helium atom

Multi-electron problem cannot be solved exactly

$$Z = 2$$

$$\left\{ -\frac{1}{2} \nabla_1^2 - \frac{1}{2} \nabla_2^2 - \frac{Z}{r_1} - \frac{Z}{r_2} + \frac{1}{r_{12}} \right\} \psi(\mathbf{r}_1, \mathbf{r}_2) = E \psi(\mathbf{r}_1, \mathbf{r}_2)$$

$$\psi(\mathbf{r}_1, \mathbf{r}_2) \neq \psi_{1s}(\mathbf{r}_1)\psi_{1s}(\mathbf{r}_2)$$

Electrons are fermions, not ideal particle

Ground state

$$\psi(1,2)_{1s^2} = \psi_{1s}(\mathbf{r}_1)\psi_{1s}(\mathbf{r}_2) \times \frac{1}{\sqrt{2}}(|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle)$$

excited state, singlet

$$\psi^+(1,2)_{1s2s} = \frac{1}{\sqrt{2}}(\psi_{1s}(\mathbf{r}_1)\psi_{2s}(\mathbf{r}_2) + \psi_{2s}(\mathbf{r}_1)\psi_{1s}(\mathbf{r}_2)) \times \frac{1}{\sqrt{2}}(|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle)$$

excited state, triplet

$$\psi^-(1,2)_{1s2s} = \frac{1}{\sqrt{2}}(\psi_{1s}(\mathbf{r}_1)\psi_{2s}(\mathbf{r}_2) - \psi_{2s}(\mathbf{r}_1)\psi_{1s}(\mathbf{r}_2)) \times \begin{pmatrix} |\uparrow\uparrow\rangle \\ \frac{1}{\sqrt{2}}(|\uparrow\downarrow\rangle + |\downarrow\uparrow\rangle) \\ |\downarrow\downarrow\rangle \end{pmatrix}$$

$$E^\pm = \underbrace{I(1s) + I(2s)}_{\text{one electron integral}} + \underbrace{J(1s,2s)}_{\text{Coulomb integral}} \pm \underbrace{K(1s,2s)}_{\text{exchange integral}}$$

Triplet has lower energy
Hund's rule

Schrödinger equation

$$i\hbar \frac{\partial}{\partial t} \psi(t, \vec{x}) = -\frac{\hbar^2}{2m} \Delta \psi(t, \vec{x}) + V(\vec{x}) \psi(t, \vec{x}) = H(\vec{x}) \psi(t, \vec{x}) \quad \psi(t, \vec{x}) = e^{-it \frac{E}{\hbar}} \psi(\vec{x})$$

1d stationary Schrödinger equation $H(\vec{x}) \psi(\vec{x}) = E \psi(\vec{x}) \quad \left(-\frac{\hbar^2}{2m} \frac{d^2}{dx^2} + V(x)\right) \psi(x) = E \psi(x)$

$$\frac{d^2 \psi(x)}{dx^2} + \frac{2m}{\hbar^2} (E - V(x)) \psi(x) = 0 \quad \frac{d^2 \psi(x)}{dx^2} + k^2(x) \psi(x) = 0 \quad \rightarrow \quad \psi''(x) = -k^2(x) \psi(x)$$

$$k^2(x) = \gamma^2 (\epsilon - \nu(x)) \quad \gamma^2 = \frac{2mV_0}{\hbar^2} \quad \epsilon = \frac{E}{V_0} \quad \nu(x) = \frac{V(x)}{V_0}$$

Numerov's algorithm

$$x = h \cdot r$$

$$\psi(r+1) = \psi(r) + h\psi'(r) + \frac{1}{2}h^2\psi''(r) + \frac{1}{6}h^3\psi^{(3)}(r) + \frac{1}{24}h^4\psi^{(4)}(r) + \dots$$

$$\psi(r-1) = \psi(r) - h\psi'(r) + \frac{1}{2}h^2\psi''(r) - \frac{1}{6}h^3\psi^{(3)}(r) + \frac{1}{24}h^4\psi^{(4)}(r) + \dots$$

$$\psi(r+1) + \psi(r-1) = 2\psi(r) + h^2\psi''(r) + \frac{1}{12}h^4\psi^{(4)}(r) + O(h^6) \quad \psi^{(4)}(r) = \frac{\psi''(r+1) + \psi''(r-1) - 2\psi''(r)}{h^2} + O(h^2)$$

$$= 2\psi(r) + h^2\psi''(r) + \frac{1}{12}h^2(\psi''(r+1) + \psi''(r-1) - 2\psi''(r)) + O(h^6)$$

$$\left(1 + \frac{h^2}{12}k^2(r+1)\right)\psi(r+1) = 2\left(1 + \frac{h^2}{12}k^2(r)\right)\psi(r) - \left(1 + \frac{h^2}{12}k^2(r-1)\right)\psi(r-1) - h^2k^2(r)\psi(r) + O(h^6)$$

$$= 2\left(1 - \frac{5}{12}h^2k^2(r)\right)\psi(r) - \left(1 + \frac{h^2}{12}k^2(r-1)\right)\psi(r-1) + O(h^6)$$

$$\psi(r+1) = \frac{2\left(1 - \frac{5}{12}h^2k^2(r)\right)\psi(r) - \left(1 + \frac{h^2}{12}k^2(r-1)\right)\psi(r-1)}{1 + \frac{1}{12}h^2k^2(r+1)}$$

given $\psi(0)$ and $\psi(1)$

Matrix Numerov's algorithm

$$\psi(r+1) + \psi(r-1) = 2\psi(r) + h^2\psi''(r) + \frac{1}{12}h^2(\psi''(r+1) + \psi''(r-1) - 2\psi''(r)) + O(h^6)$$

$$\psi''(x) = -k^2(x)\psi(x)$$

$$\psi(r+1) - 2\psi(r) + \psi(r-1) = -\frac{h^2}{12}k^2(r+1)\psi(r+1) - \frac{10h^2}{12}k^2(r)\psi(r) - \frac{h^2}{12}k^2(r-1)\psi(r-1)$$

$$k^2(x) = \gamma^2(\epsilon - \nu(x)) \quad \gamma^2 = \frac{2mV_0}{\hbar^2} \quad \epsilon = \frac{E}{V_0} \quad \nu(x) = \frac{V(x)}{V_0}$$

$$-\frac{1}{\gamma^2} \left(\frac{\psi(r+1) - 2\psi(r) + \psi(r-1)}{h^2} \right) + \frac{\nu(r+1)\psi(r+1) + 10\nu(r)\psi(r) + \nu(r-1)\psi(r-1)}{12} = \epsilon \frac{\psi(r+1) + 10\psi(r) + \psi(r-1)}{12}$$

$$\Psi = \begin{pmatrix} \psi(1) \\ \psi(2) \\ \psi(3) \\ \vdots \\ \psi(N-1) \\ \psi(N) \end{pmatrix}$$

$$\mathbf{A} = \frac{1}{h^2}(\mathbb{1}_{-1} - 2\mathbb{1}_0 + \mathbb{1}_1)$$

$$\mathbf{B} = \frac{1}{12}(\mathbb{1}_{-1} + 10\mathbb{1}_0 + \mathbb{1}_1)$$

$$\nu = \begin{pmatrix} \nu_1 & 0 & \dots & 0 & 0 \\ 0 & \nu_2 & \dots & 0 & 0 \\ \vdots & 0 & \ddots & \vdots & \\ 0 & 0 & \dots & \nu_{N-1} & 0 \\ 0 & 0 & \dots & 0 & \nu_N \end{pmatrix}$$

$$\left(-\frac{1}{\gamma^2} \mathbf{B}^{-1} \mathbf{A} + \nu \right) \Psi = \epsilon \Psi$$

Hamiltonian matrix

boundary condition $\psi(1)$ has no left neighbour

boundary condition $\psi(N)$ has no right neighbour

Solution for square well

$$\Psi'' + k^2\Psi = 0 \quad k^2 = \gamma^2(\epsilon - \nu)$$

$$\nu(0) = \nu(L) = \infty$$


$$\frac{2m}{\hbar^2} = 1, \gamma^2 = 1 \quad \nu(x) = 0, x \in (0, L)$$

$$\psi_n(x) = \sqrt{2} \sin(k_n x) = \sqrt{2} \sin(n\pi x) \quad k_n = n\pi/L \quad \psi(0) = \psi(L) = 0$$


$$E_n = \frac{n^2 \pi^2}{L^2 \gamma^2} \quad n = 1, 2, 3, \dots$$

$n = 0, \psi_n(x) = 0$ trivial solution, no particle

https://en.wikipedia.org/wiki/Particle_in_a_box



Solution for harmonic potential



$$-\frac{\hbar^2}{2m} \frac{d^2}{dx^2} \psi + \frac{1}{2} m \omega^2 x^2 \psi = E \psi \quad \nu(x) = \frac{1}{2} m \omega^2 x^2$$

$$-\frac{1}{2} \frac{d^2}{dx^2} \psi + \frac{1}{2} \left(\frac{m\omega}{\hbar}\right)^2 x^2 \psi = \frac{m\omega}{\hbar} \frac{E}{\hbar\omega} \psi \quad \frac{m\omega}{\hbar} = 1 \quad \epsilon = \frac{E}{\hbar\omega}$$

$$\left(-\frac{1}{2} \mathbf{B}^{-1} \mathbf{A} + \nu\right) \Psi = \epsilon \Psi \quad \nu = \frac{1}{2} x^2$$

$$\psi_n(x) = \frac{1}{\sqrt{2^n n!}} \left(\frac{m\omega}{\pi\hbar}\right)^{1/4} e^{-\frac{m\omega x^2}{2\hbar}} H_n\left(\sqrt{\frac{m\omega}{\hbar}} x\right) \quad n = 0, 1, 2, \dots$$

$$E_n = \hbar\omega(n + 1/2)$$

https://en.wikipedia.org/wiki/Quantum_harmonic_oscillator

Time-dependent Schrödinger equation

$$i\hbar \frac{\partial}{\partial t} \psi(t, \vec{x}) = -\frac{\hbar^2}{2m} \Delta \psi(t, \vec{x}) + V(\vec{x}) \psi(t, \vec{x}) = H(\vec{x}) \psi(t, \vec{x}) \quad \psi(t, x) \rightarrow \psi(t_n, x_r) = \psi_r^n$$

FTCS scheme

$$i\hbar \frac{\psi_r^{n+1} - \psi_r^n}{\tau} = -\frac{\hbar^2}{2m} \frac{\psi_{r+1}^n + \psi_{r-1}^n - 2\psi_r^n}{h^2} + V(r) \psi_r^n = \sum_{s=1}^N H_{r,s} \psi_s^n \quad N \times N$$

$$\vec{\psi}^n = \begin{pmatrix} \psi_0^n \\ \psi_1^n \\ \psi_2^n \\ \vdots \\ \psi_{N-1}^n \end{pmatrix}$$

$$H_{r,s} = -\frac{\hbar^2}{2m} \frac{\delta_{r+1,s} + \delta_{r-1,s} - 2\delta_{r,s}}{h^2} + V(r) \delta_{r,s}$$

von Neumann stability analysis $\psi_r^n = A^n e^{ikrh}$

$$\vec{\psi}^{n+1} = \left(1 - \frac{i\tau}{\hbar} H\right) \vec{\psi}^n \quad \text{for the free case } V(x) = 0$$

$$A = 1 - i \frac{\hbar\tau}{mh^2} (\cos(kh) - 1) \quad |A|^2 > 1$$

Fully-Implicit scheme

$$i\hbar \frac{\psi_r^{n+1} - \psi_r^n}{\tau} = \sum_{s=1}^N H_{r,s} \psi_s^{n+1}$$

$$\vec{\psi}^{n+1} = \vec{\psi}^n - \frac{i\tau}{\hbar} H \vec{\psi}^{n+1} \quad \left(1 + \frac{i\tau}{\hbar} H\right) \vec{\psi}^{n+1} = \vec{\psi}^n \quad \vec{\psi}^{n+1} = \left(1 + \frac{i\tau}{\hbar} H\right)^{-1} \vec{\psi}^n$$

von Neumann stability analysis $\psi_r^n = A^n e^{ikrh}$

for the free case

$$V(x) = 0$$

$$A = \frac{1}{1 + i \frac{\hbar\tau}{mh^2} (\cos(kh) - 1)} \quad |A| < 1$$

do not preserve the norm of wave function, not unitary

$\langle \psi(0) | \psi(0) \rangle = 1$ $\psi(t) = e^{-itH/\hbar} \psi(0)$ preserve the normalisation H is an hermitian matrix
 $H = H^\dagger$

$(e^{-itH/\hbar})^\dagger = e^{itH/\hbar} = (e^{-itH/\hbar})^{-1}$ $\langle \psi(t) | \psi(t) \rangle = \langle \psi(0) | \underbrace{e^{itH/\hbar} \cdot e^{-itH/\hbar}}_{=1} | \psi(0) \rangle$

FTCS scheme $e^{-i\tau H/\hbar} \approx (1 - \frac{i\tau}{\hbar} H)$ breaks the unitarity

Fully-Implicit scheme $e^{-i\tau H/\hbar} \approx (1 + \frac{i\tau}{\hbar} H)^{-1}$

$(e^{-ix})^\dagger = e^{ix} = (e^{-ix})^{-1}$ but $(1 \pm ix)^\dagger = 1 \mp ix \neq 1/(1 \pm ix)$

Crank-Nicolson scheme $i\hbar \frac{\psi_r^{n+1} - \psi_r^n}{\tau} = \sum_{s=1}^N H_{r,s} \frac{1}{2} (\underbrace{\psi_s^n}_{FTCS} + \underbrace{\psi_s^{n+1}}_{Fully-implicit})$

unitary discretisation scheme

$(1 + \frac{i\tau}{2\hbar} H) \vec{\psi}^{n+1} = (1 - \frac{i\tau}{2\hbar} H) \vec{\psi}^n$ $\vec{\psi}^{n+1} = \vec{\psi}^n - \frac{i\tau}{2\hbar} H(\vec{\psi}^n + \vec{\psi}^{n+1})$



$\vec{\psi}^{n+1} = \underbrace{(1 + \frac{i\tau}{2\hbar} H)^{-1} (1 - \frac{i\tau}{2\hbar} H)}_{\text{unitary}} \vec{\psi}^n$ $(\frac{1 - ix}{1 + ix})^\dagger = \frac{1 + ix}{1 - ix} = (\frac{1 - ix}{1 + ix})^{-1}$

Chap.20.2.

unitary

Can you see it? $|\frac{1 - ix}{1 + ix}| = 1$

Operator splitting

$$H = -\frac{\hbar^2}{2m}(\Delta_x + \Delta_y + \Delta_z + V(\vec{x})) = H_{kin} + H_{pot}$$

$$\psi(t + \tau) = U(\tau)\psi(t) = e^{-i\frac{\tau}{\hbar}H}\psi(t) \quad e^{-i\frac{\tau}{\hbar}H}|\psi\rangle \rightarrow e^{-i\frac{\tau}{\hbar}H_{kin}}e^{-i\frac{\tau}{\hbar}H_{pot}}|\psi\rangle$$

Trotter-Suzuki decomposition $e^{\tau(A+B)} = e^{\tau A} \cdot e^{\tau B} + O(\tau^2)$ in the eigen basis $e^{-i\frac{\tau}{\hbar}V(\vec{x})}\psi(\vec{x})$

Baker-Campbell-Hausdorff $e^X e^Y = e^{X+Y+\frac{1}{2}[X,Y]+\frac{1}{12}([X,[X,Y]]-[Y,[X,Y]])+\dots}$ $e^{-i\frac{\tau}{\hbar}\frac{\hbar^2 k_x^2}{2m}}\psi(k_x)$

But the Eigen bases for H_{kin} and H_{pot} are different

Spectral methods $i\hbar\frac{\partial}{\partial t}\vec{\Psi}(t) = H\vec{\Psi}(t) \quad H\vec{\Psi} = E\vec{\Psi} \quad \vec{\Psi}(t) = e^{-itE/\hbar}\vec{\Psi}(0)$

If one knows all the eigenvalues E_n and corresponding eigenstates $\vec{\phi}_n$ of the H

$$\vec{\Psi}(t=0) = \sum_n c_n \vec{\phi}_n$$

$$\vec{\Psi}(t) = \sum_n c_n e^{-i\frac{t}{\hbar}E_n} \vec{\phi}_n$$

All these point to the diagonalization of the Hamiltonian matrix